

Multi-Pronged Attack Against Arthritis

References

1. Danelich IM, Wright SS, Lose JM, et al. Safety of nonsteroidal antiinflammatory drugs in patients with cardiovascular disease. *Pharmacotherapy*. 2015;35(5):520-35.
2. Lanas A, Boers M, Nuevo J. Gastrointestinal events in at-risk patients starting non-steroidal anti-inflammatory drugs (NSAIDs) for rheumatic diseases: the EVIDENCE study of European routine practice. *Ann Rheum Dis*. 2015;74(4):675-81.
3. Park KK, Choi CH, Ha CW, et al. The Effects of Adherence to Non-Steroidal Anti-Inflammatory Drugs and Factors Influencing Drug Adherence in Patients with Knee Osteoarthritis. *J Korean Med Sci*. 2016;31(5):795-800.
4. Santilli F, Boccatonda A, Davi G, et al. The Coxib case: Are EP receptors really guilty? *Atherosclerosis*. 2016;249:164-73.
5. Singh DP, Borse SP, Nivsarkar M. Clinical importance of nonsteroidal anti-inflammatory drug enteropathy: the relevance of tumor necrosis factor as a promising target. *Transl Res*. 2016.
6. Tonge DP, Pearson MJ, Jones SW. The hallmarks of osteoarthritis and the potential to develop personalised disease-modifying pharmacological therapeutics. *Osteoarthritis Cartilage*. 2014;22(5):609-21.
7. Sampalis JS, Brownell LA. A randomized, double blind, placebo and active comparator controlled pilot study of UP446, a novel dual pathway inhibitor anti-inflammatory agent of botanical origin. *Nutr J*. 2012;11:21.
8. Arjmandi BH, Ormsbee LT, Elam ML, et al. A combination of Scutellaria baicalensis and Acacia catechu extracts for short-term symptomatic relief of joint discomfort associated with osteoarthritis of the knee. *J Med Food*. 2014;17(6):707-13.
9. Unigen Inc. Amlexin: A New Approach to Joint Protection. Unpublished data. 2015.
10. Saberi Hosnijeh F, Siebuhr AS, Uitterlinden AG, et al. Association between biomarkers of tissue inflammation and progression of osteoarthritis: evidence from the Rotterdam study cohort. *Arthritis Res Ther*. 2015;18:81.
11. Garner P, Piperno M, Gineyts E, et al. Cross sectional evaluation of biochemical markers of bone, cartilage, and synovial tissue metabolism in patients with knee osteoarthritis: relations with disease activity and joint damage. *Ann Rheum Dis*. 2001;60(6):619-26.
12. Sokolove J, Lepus CM. Role of inflammation in the pathogenesis of osteoarthritis: latest findings and interpretations. *Ther Adv Musculoskelet Dis*. 2013;5(2):77-94.
13. Yimam M, Brownell L, Pantier M, et al. UP446, analgesic and anti-inflammatory botanical composition. *Pharmacognosy Res*. 2013;5(3):139-45.
14. Tseng-Crank J, Sung S, Jia Q, et al. A medicinal plant extract of Scutellaria Baicalensis and Acacia catechu reduced LPS-stimulated gene expression in immune cells: a comprehensive genomic study using QPCR, ELISA, and microarray. *J Diet Suppl*. 2010;7(3):253-72.
15. Morgan SL, Baggott JE, Moreland L, et al. The safety of flavocoxid, a medical food, in the dietary management of knee osteoarthritis. *J Med Food*. 2009;12(5):1143-8.
16. Burnett BP, Jia Q, Zhao Y, et al. A medicinal extract of Scutellaria baicalensis and Acacia catechu acts as a dual inhibitor of cyclooxygenase and 5-lipoxygenase to reduce inflammation. *J Med Food*. 2007;10(3):442-51.
17. Burnett BP, Levy RM. 5-Lipoxygenase metabolic contributions to NSAID-induced organ toxicity. *Adv Ther*. 2012;29(2):79-98.